Neuroscience Letters, 85 (1988) 375–379
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Agnati, L.F., see Fuxe, K., 163
Amann, R., Hankins, M.W. and Dray, A., Actions of neuropeptide K and calcitonin gene-related peptide on inferior mesenteric ganglion cells–tachykinin interactions with non-cholinergic potentials evoked by ureteric nerve stimulation, 125
Aquilonius, S.-M., see d'Argy, R., 304
Arango, V., see Ernsberger, P., 179
Armengaud, C., Leitner, L.-M., Malber, C.-H., Roumy, M., Ruckebusch, M. and Sutra, J.F., Comparison of the monoamine and catabolite content in the cat and rabbit carotid bodies, 153
Aston-Jones, G., see Pieribone, V.A., 297

Barker, J.L., see Harrison, N.L., 105
Beadle, D.J., see Horseman, B.G., 65
Beart, P.M., Nicolopoulos, L.S., West, D.C. and Headley, P.M., An excitatory amino acid projection from ventromedial hypothalamus to periaqueductal gray in the rat: autoradiographic and electrophysiological evidence, 205
Bellini, F., see Pantaleo, T., 56
Benavidès, J., see Savasta, M., 119
Bennett, T., see Harland, D., 239
Bergström, M., see d'Argy, R., 304
Bermudez, I., see Horseman, B.G., 65
Bock, S.E., see Deaton, M.A., 267
Bohn, M.C., see Pieribone, V.A., 297
Bolin, L.M., Thomas, A.R., Mayer, D.N. and Rouse, R.V., Independent developmental regulation of migratory granule neurons and their cerebellar ligand in the mouse, 158
Bonshek, R.E., see Mann, D.M.A., 277
Boris-Möller, F., Drakenberg, T., Elmdén, K., Forsén, S. and Siesjö, B.K., Evidence against major compartmentalization of H^+ in ischemic rat brain tissue, 113
Bose, R., see Hall, A., 110
Bosler, O., see Guy, J., 9
Brachet, P., see Dicou, E., 19
Bruno, J.P., see Weihmuller, F.B., 137
Bulant, M., see Liao, N., 24
Burden, H.W., see Klein, C.M., 217

Chapman, J.A., see Murphy, J.A., 255
Choi, D.W., see Wong, B.Y., 261
Clarke, I.J., see Thomas, G.B., 143
Convents, A., see De Keyser, J., 272
Coulter, D.A., see Wong, B.Y., 261
Cummins, J.T., see Thomas, G.B., 143

d'Argy, R., Gillberg, P.-G., Stålnacke, C.-G., Persson, A., Bergström, M., Långström, B., Schoeps, K.-O. and Aquilonius, S.-M., In vivo and in vitro receptor autoradiography of the human brain using an ^{11}C-labelled benzodiazepine analogue, 304
Davis, H.L., Desypris, G. and Lui, S.L., Comparison of fiber size and phenotypic gene expression in muscles of dystrophic C57BL/6J DY2J/DY2J mice, 370
Deaton, M.A., Bock, S.E., Zwick, H. and Freeman, J.A., Common properties shared by growth-associated proteins of the regenerating optic nerve of goldfish *(C. auratus)*, 267
De Backer, J.-P., see De Keyser, J., 272
De Keyser, J., Ebinger, G., De Backer, J.-P., Convents, A., Vanderheyden, P. and Vauquelin, G., Subtypes of adrenergic and dopaminergic receptors in bovine cerebral blood vessels, 272
Despres, G., Giry, N. and Romand, R., Immunohistochemical localisation of nerve growth factor-like protein in the organ of Corti of the developing rat, 5
Desypris, G., see Davis, H.L., 370
Dicou, E., Lee, J. and Brachet, P., Co-localization of the nerve growth factor precursor protein and mRNA in the mouse submandibular gland, 19
Drakenberg, T., see Boris-Möller, F., 113
Drapeau, P., Effects of kainic acid lesions on calcium uptake and dopamine release in nerve endings isolated from rat striata, 233
Dray, A., see Amann, R., 125
Dubois, A., see Savasta, M., 119
Duranti, R., see Pantaleo, T., 56
Dutar, P. and Nicoll, R.A., Stimulation of phosphatidylinositol (PI) turnover may

mediate the muscarinic suppression of the M-current in hippocampal pyramidal cell, 89

Ebinger, G., see De Keyser, J., 272
Eisenman, L.M., see Heckroth, J.A., 199
Elde, R., see Hill, E.L., 172
Elmdén, K., see Boris-Möller, F., 113
Ernsberger, P., Arango, V. and Reis, D.J., A high density of muscarinic receptors in the rostral ventrolateral medulla of the rat is revealed by correction for autoradiographic efficiency, 179

Fan, L., see Shu, S., 169
Feeser, H.R., see Prince, D.A., 291
Forsén, S., see Boris-Möller, F., 113
Freeman, J.A., see Deaton, M.A., 267
Fuxe, K., von Euler, G., Agnati, L.F. and Ögren, S.-O., Galanin selectively modulates 5-hydroxytryptamine 1A receptors in the rat ventral limbic cortex, 163
Fuxe, K., see Ögren, S.O., 333

Gardiner, S.M., see Harland, D., 239
Ghiani, P., Uva, B., Vallarino, M., Mandich, A. and Masini, M.A., Angiotensin II specific receptors in subcommissural organ, 212
Gibbens, H., see Parsons, C.G., 322
Gillberg, P.-G., see d'Argy, R., 304
Giménez-Amaya, J.M., Afferent connections to the pontine nuclei from the cortex of the anterior ectosylvian sulcus in the cat, 29
Giry, N., see Despres, G., 5
Glavin, G., see Hall, A., 110
Graeber, M.B., Tetzlaff, W., Streit, W.J. and Kreutzberg, G.W., Microglial cells but not astrocytes undergo mitosis following rat facial nerve axotomy, 317
Griffioen, A.W, see Witter, M.P., 193
Gustafsson, B., Huang, Y.-Y. and Wigström, H., Phorbol ester-induced synaptic potentiation differs from long-term potentiation in the guinea pig hippocampus in vitro, 77
Guy, J., Pelletier, G. and Bosler, O., Serotonin innervation of neuropeptide Y-containing neurons in the rat arcuate nucleus, 9

Hadjiconstantinou, M., see Weihmuller, F.B., 137
Hall, A., Bose, R., Glavin, G. and Pinsky, C., Effects of 4-acetylpyridine on stress ulcer formation in mice, 110
Hankins, M.W., see Amann, R., 125
Hansen, S., see Perry, T.L., 101
Harada, M., see Ozaki, N., 228
Harland, D., Bennett, T. and Gardiner, S.M., Cardiovascular actions of neuropeptide Y in the hypothalamic paraventricular nucleus of conscious Long Evans and Brattleboro rats, 239
Harrison, N.L., Lange, G.D. and Barker, J.L., (–)-Baclofen activates presynaptic $GABA_B$ receptors on GABAergic inhibitory neurons from embryonic rat hippocampus, 105
Headley, P.M., see Beart, P.M., 205
Headley, P.M., see Parsons, C.G., 322
Heckroth, J.A. and Eisenman, L.M., The olivocerebellar projection in 'lurcher' mutant mice, 199
Helkala, E.-L., see Soininen, H., 131
Herz, A., see Weihe, E., 187
Hill, E.L. and Elde, R., Calcitonin gene-related peptide-immunoreactive nerve fibers in mandibular periosteum of rat: evidence for primary afferent origin, 172
Horseman, B.G., Seymour, C., Bermudez, I. and Beadle, D.J., The effects of L-glutamate on cultured insect neurones, 65
Hoyer, D., Waeber, C., Pazos, A., Probst, A. and Palacios, J.M., Identification of a 5-HT_1 recognition site in human brain membranes different from 5-HT_{1B} and 5-HT_{1C} sites, 357
Huang, Y.-Y., see Gustafsson, B., 77
Huidobro-Toro, J.P. and Parada, S., Co-transmission in the rat vas deferens: postjunctional synergism of noradrenaline and adenosine 5′-triphosphate, 339

Iijima, T., see Kudo, Y., 345
Ikeda, K. and Tomonaga, M., Creatine kinase immunoreactivity: localization in nerve terminals in the hypothalamic area and superior colliculus of the mouse brain, 51
Imeri, L., see Marini, G., 223
Imura, H., see Inoue, T., 363
Imura, H., see Inoue, T., 95
Inoue, T., Kato, Y., Koshiyama, H., Yanaihara, N. and Imura, H., Interaction between vasoactive intestinal polypeptide (VIP) and peptide histidine isoleucine (PHI) in stimulating the secretion of prolactin from rat anterior pituitary cells in vitro, 363
Inoue, T., Kato, Y., Koshiyama, H., Yanaihara, N. and Imura, H., Galanin stimulates the release of vasoactive intestinal

polypeptide from perifused hypothalamic fragments in vitro and from periventricular structures into the cerebrospinal fluid in vivo in the rat, 95
Itzhak, Y. and Khouri, M., Regulation of the binding of σ- and phencyclidine (PCP)-receptor ligands in rat brain membranes by guanine nucleotides and ions, 147
Iwai, C., see Ochiai, H., 14

Jolkkonen, J., see Soininen, H., 131
Jones, K., see Perry, T.L., 101
Jorritsma-Byham, B., see Witter, M.P., 193
Ju, G., see Shu, S., 169

Kaneda, N., see Ozaki, N., 228
Kasahara, Y., see Ozaki, N., 228
Kato, Y., see Inoue, T., 363
Kato, Y., see Inoue, T., 95
Khouri, M., see Itzhak, Y., 147
Kiuchi, K., see Ozaki, N., 228
Klein, C.M. and Burden, H.W., Anatomical localization of afferent and postganglionic sympathetic neurons innervating the rat ovary, 217
Koshiyama, H., see Inoue, T., 363
Koshiyama, H., see Inoue, T., 95
Kosinski, R.J., Lee, H.S. and Mihailoff, G.A., A double retrograde fluorescent tracing analysis of dorsal column nuclear projections to the basilar pontine nuclei, thalamus, and superior colliculus in the rat, 40
Kreutzberg, G.W., see Graeber, M.B., 317
Krijnen, J.L.M., see Witter, M.P., 193
Krishtal, O.A., Smirnov, S.V. and Osipchuk, Yu.V., Changes in the state of the excitatory synaptic system in the hippocampus on prolonged exposure to excitatory amino acids and antagonists, 82
Kudo, M., see Tokuno, H., 1
Kudo, Y., Ogura, A. and Iijima, T., Stimulation of muscarinic receptor in hippocampal neuron induces characteristic increase in cytosolic free Ca^{2+} concentration, 345

Lange, G.D., see Harrison, N.L., 105
Långström, B., see d'Argy, R., 304
Laulumaa, V., see Soininen, H., 131
Lauweryns, J.M. and Van Ranst, L., Protein gene product 9.5 expression in the lungs of humans and other mammals. Immunocytochemical detection in neuroepithelial bodies, neuroendocrine cells and nerves, 311
Lee, H.S., see Kosinski, R.J., 40
Lee, J., see Dicou, E., 19
Leibold, A., see Weihe, E., 187
Leitner, L.-M., see Armengaud, C., 153
Liao, N., Bulant, M., Nicolas, P., Vaudry, H. and Pelletier, G., Immunocytochemical distribution of neurons containing a peptide derived from thyrotropin-releasing hormone precursor in the rat brain, 24
Liguri, G., Nassi, P., Taddei, N., Nediani, C. and Ramponi, G., Post-mortem modifications of the specific activity of some brain enzymes, 244
Lui, S.L., see Davis, H.L., 370
Lynn, B. and Shakhanbeh, J., Properties of Aδ high threshold mechanoreceptors in the rat hairy and glabrous skin and their response to heat, 71

Maeyama, K., see Nakamura, S., 329
Magnago, T.S.I., see Parsons, C.G., 322
Malber, C.-H., see Armengaud, C., 153
Mancia, M., see Marini, G., 223
Mandich, A., see Ghiani, P., 212
Mann, D.M.A., Bonshek, R.E, Marcyniuk, B., Stoddart, R.W. and Torgerson, E., Saccharides of senile plaques and neurofibrillary tangles in Alzheimer's disease, 277
Marcyniuk, B., see Mann, D.M.A., 277
Marini, G., Imeri, L. and Mancia, M., Changes in sleep–waking cycle induced by lesions of medialis dorsalis thalamic nuclei in the cat, 223
Martinez, S., see Puelles, L., 35
Martinez-de-la-Torre, M., see Puelles, L., 35
Masini, M.A., see Ghiani, P., 212
Mayer, D.N., see Bolin, L.M., 158
McDonald, A.J., Projections of the intermediate subdivision of the central amygdaloid nucleus to the bed nucleus of the stria terminalis and medial diencephalon, 285
Mihailoff, G.A., see Kosinski, R.J., 40
Millan, M.J., see Weihe, E., 187
Miura, Y., see Ozaki, N., 228
Mogi, M., see Ozaki, N., 228
Moriizumi, T., see Tokuno, H., 1
Murphy, J.A., Chapman, J.A., Suckling, A.J. and Rumsby, M.G., Protein kinase C activity in soluble fractions from glial cells in primary culture and subcultures, 255

Nagatsu, T., see Ozaki, N., 228
Nakahara, D., see Ozaki, N., 228

Nakai, Y., see Ochiai, H., 14
Nakamura, S., Sakaguchi, T., Shirokawa, T., Yamatodani, A., Maeyama, K. and Wada, H., Changes in the electrical activity of locus coeruleus neurons in pregnant rats, 329
Nakamura, Y., see Tokuno, H., 1
Nassi, P., see Liguri, G., 244
Nediani, C., see Liguri, G., 244
Nicolas, P., see Liao, N., 24
Nicoll, R.A., see Dutar, P., 89
Nicolopoulos, L.S., see Beart, P.M., 205
Nishizaki, T., Yamauchi, R. and Okada, Y., Enhancement of the oxygen consumption in the hippocampal slices of the guinea pig induced by glutamate and its related substances, 61
Nistri, A., see Sivilotti, L., 249
Nohr, D., see Weihe, E., 187

Ochiai, H., Iwai, C. and Nakai, Y., Ultrastructural demonstration of the catecholaminergic innervation of vasopressin neurons in the paraventricular nucleus of the rat by double-labeling immunocytochemistry, 14
Ögren, S.-O., see Fuxe, K., 163
Ögren, S.O. and Fuxe, K., D_1- and D_2-receptor antagonist induce catalepsy via different efferent. Striatal pathways, 333
Ogura, A., see Kudo, Y., 345
Okada, Y., see Nishizaki, T., 61
Osipchuk, Yu.V., see Krishtal, O.A., 82
Ozaki, N., Nakahara, D., Mogi, M., Harada, M., Kiuchi, K., Kaneda, N., Miura, Y., Kasahara, Y. and Nagatsu, T., Inactivation of tyrosine hydroxylase in rat striatum by 1-methyl-4-phenylpyridinium ion (MMP^+), 228

Palacios, J.M., see Hoyer, D., 357
Pantaleo, T., Duranti, R. and Bellini, F., Effects of heterotopic ischemic pain on muscular pain threshold and blink reflex in humans, 56
Parada, S., see Huidobro-Toro, J.P., 339
Parsons, C.G., Gibbens, H., Magnago, T.S.I. and Headley, P.M., At which 'sigma' site are the spinal actions of ketamine mediated?, 322
Partanen, J., see Soininen, H., 131
Pazos, A., see Hoyer, D., 357
Pelletier, G., see Guy, J., 9
Pelletier, G., see Liao, N., 24
Perry, T.L., Jones, K. and Hansen, S., Tetrahydroisoquinoline lacks dopaminergic nigrostriatal neurotoxicity in mice, 101
Persson, A., see d'Argy, R., 304
Pieribone, V.A., Aston-Jones, G. and Bohn, M.C., Adrenergic and non-adrenergic neurons in the C1 and C3 areas project to locus coeruleus: a fluorescent double labeling study, 297
Pinsky, C., see Hall, A., 110
Prince, D.A. and Feeser, H.R., Dextromethorphan protects against cerebral infarction in a rat model of hypoxia-ischemia, 291
Prince, D.A., see Wong, B.Y., 261
Probst, A., see Hoyer, D., 357
Puelles, L., Martinez, S. and Martinez-de-la-Torre, M., The locus of optic nerve head representation in the retinotopic projection over nucleus geniculatus lateralis ventralis and nucleus griseum tectalis in the chick also lacks a retinal projection, 35

Radhakishun, F.S., Van Ree, J.M. and Westerink, B.H.C., Scheduled eating increases dopamine release in the nucleus accumbens of foor-deprived rats as assessed with on-line brain dialysis, 351
Ramponi, G., see Liguri, G., 244
Reinikainen, K., see Soininen, H., 131
Reis, D.J., see Ernsberger, P., 179
Riekkinen, P.J., see Soininen, H., 131
Romand, R., see Despres, G., 5
Roumy, M., see Armengaud, C., 153
Rouse, R.V., see Bolin, L.M., 158
Ruckebusch, M., see Armengaud, C., 153
Rumsby, M.G., see Murphy, J.A., 255

Sakaguchi, T., see Nakamura, S., 329
Savasta, M., Dubois, A., Benavidès, J. and Scatton, B., Different plasticity changes in D_1 and D_2 receptors in rat striatal subregions following impairment of dopaminergic transmission, 119
Scatton, B., see Savasta, M., 119
Schoeps, K.-O., see d'Argy, R., 304
Seymour, C., see Horseman, B.G., 65
Shakhanbeh, J., see Lynn, B., 71
Shirokawa, T., see Nakamura, S., 329
Shu, S., Ju, G. and Fan, L., The glucose oxidase–DAB–nickel method in peroxidase histochemistry of the nervous system, 169
Siesjö, B.K., see Boris-Möller, F., 113

Sivilotti, L. and Nistri, A., Complex effects of baclofen on synaptic transmission of the frog optic tectum in vitro, 249
Smirnov, S.V., see Krishtal, O.A., 82
Soininen, H., Riekkinen, P.J., Partanen, J., Helkala, E.-L., Laulumaa, V., Jolkkonen, J. and Reinikainen, K., Cerebrospinal fluid somatostatin correlates with spectral EEG variables and with parietotemporal cognitive dysfunction in Alzheimer patients, 131
Stålnacke, C.-G., see d'Argy, R., 304
Stoddart, R.W., see Mann, D.M.A., 277
Streit, W.J., see Graeber, M.B., 317
Suckling, A.J., see Murphy, J.A., 255
Sutra, J.F., see Armengaud, C., 153

Taddei, N., see Liguri, G., 244
Tetzlaff, W., see Graeber, M.B., 317
Thomas, A.R., see Bolin, L.M., 158
Thomas, G.B., Cummins, J.T. and Clarke, I.J., Secretion of prolactin in response to serotonin requires an intact hypothalamo-pituitary axis in the ewe, 143
Tokuno, H., Moriizumi, T., Kudo, M. and Nakamura, Y., A morphological evidence for monosynaptic projections from the nucleus tegmenti pedunculopontinus pars compacta (TPC) to nigrostriatal projection neurons, 1
Tomonaga, M., see Ikeda, K., 51
Torgerson, E., see Mann, D.M.A., 277

Uva, B., see Ghiani, P., 212

Vallarino, M., see Ghiani, P., 212
Vanderheyden, P., see De Keyser, J., 272
Van Ranst, L., see Lauweryns, J.M., 311
Van Ree, J.M., see Radhakishun, F.S., 351
Vaudry, H., see Liao, N., 24
Vauquelin, G., see De Keyser, J., 272
Von Euler, G., see Fuxe, K., 163

Wada, H., see Nakamura, S., 329
Waeber, C., see Hoyer, D., 357
Walz, B., Brain but not retinal glial cells have carbonic anhydrase activity in the honeybee drone, 47
Weihe, E., Millan, M.J., Leibold, A., Nohr, D. and Herz, A., Co-localization of proenkephalin- and prodynorphin-derived opioid peptides in laminae IV/V spinal neurons revealed in arthritic rats, 187
Weihmuller, F.B., Hadjiconstantinou, M. and Bruno, J.P., Acute stress or neuroleptics elicit sensorimotor deficits in MPTP-treated mice, 137
West, D.C., see Bert, P.M., 205
Westerink, B.H.C., see Radhakishun, F.S., 351
Wigström, H., see Gustafsson, B., 77
Witter, M.P., Griffioen, A.W., Jorritsma-Byham, B. and Krijnen, J.L.M., Entorhinal projections to the hippocampal CA1 region in the rat: an underestimated pathway, 193
Wong, B.Y., Coulter, D.A., Choi, D.W. and Prince, D.A., Dextrorphan and dextromethorphan, common antitussives, are antiepileptic and antagonize *N*-methyl-D-aspartate in brain slices, 261

Yamatodani, A., see Nakamura, S., 329
Yamauchi, R., see Nishizaki, T., 61
Yanaihara, N., see Inoue, T., 363
Yanaihara, N., see Inoue, T., 95

Zwick, H., see Deaton, M.A., 267